

BIRDS OF WASHINGTON

Dennis Paulson

Last revision 14 August 2021

This list includes all the bird species regularly (annually) reported in Washington state and any that have bred in the last decade. Species recorded from the state at least once but not annually are listed in a supplementary list of "Irregular Species" at the end. Common and scientific names and species order from AOU Check-list of North American Birds (Seventh Edition) and supplements through July 2021.

KEY TO STATUS OF SPECIES

PR - permanent resident (breeds unless otherwise indicated)

SR - summer resident (breeds)

SV - summer visitor (does not breed)

WV - winter visitor

M - migrant (spring & fall)

Abund. - abundant (dependably seen in large numbers)

Com. - common (dependably seen in moderate numbers)

Uncom. - uncommon (often but not always seen in small numbers)

Rare (present every year but seldom seen)

Order ANSERIFORMES

ANATIDAE (waterfowl)

Emperor Goose, *Anser canagicus*. Rare WV, mostly in coastal areas. Breeds in Alaska.

Snow Goose, *Anser caerulescens*. Large numbers winter in coastal marshes and fields in Skagit/Snohomish counties, widely increasing and now locally common in Columbia Basin. M or WV elsewhere. Breeds in Arctic (our birds mostly from Siberia).

Ross's Goose, *Anser rossii*. Rare M anywhere in state. Breeds in Canadian Arctic.

Greater White-fronted Goose, *Anser albifrons*. Com. M on outer coast, rarely elsewhere; uncom. WV west of Cascades. Breeds in Arctic.

Brant, *Branta bernicla*. Locally abund. WV and widely com. spring M on salt water, rare M east. Breeds in Arctic.

Cackling Goose, *Branta hutchinsii*. Cackling Goose, *Branta hutchinsii*. Com. WV southwest, com. M along coast, uncommon M east. Increasing dramatically. Breeds in Arctic and Subarctic.

Canada Goose, *Branta canadensis*. Com. PR (populations west of Cascades introduced); abund. M and locally com WV on coast and east of Cascades (from Arctic-breeding populations).

Trumpeter Swan, *Cygnus buccinator*. Locally com. WV on freshwater lakes and agriculture lands west of Cascades, rare east but increasing everywhere. Breeds in Alaska/BC boreal forest.

Tundra Swan, *Cygnus columbianus*. Locally com. WV on salt and fresh water and fields in western lowlands; locally com. M east of Cascades. Breeds in Arctic.

Wood Duck, *Aix sponsa*. Uncom. SR on wooded lakes; winters locally in increasing numbers where provisioned by humans.

Blue-winged Teal, *Spatula discors*. Uncom. SR in marshes, more common east.

Cinnamon Teal, *Spatula cyanoptera*. SR in marshes, com. east, locally com. west.

Northern Shoveler, *Spatula clypeata*. Com. WV, more on fresh than salt water; SR in marshes, com. east, very local west.

Gadwall, *Mareca strepera*. Com. PR in marshes, lakes and bays.

- Eurasian Wigeon, *Mareca penelope*. Uncom. WV on fresh and salt water, mostly west. Breeds in Siberian boreal forest.
- American Wigeon, *Mareca americana*. Abund. WV on fresh and salt water; uncom. SR in marshes east of Cascades, very local west.
- Mallard, *Anas platyrhynchos*. Abund. PR, more common on fresh than salt water.
- Northern Pintail, *Anas acuta*. Com. WV on coast, less common on fresh water; uncom. SR in marshes east of Cascades.
- Green-winged Teal, *Anas crecca*. Abund. WV on coast, less common on fresh water; uncom. SR in marshes east of Cascades, few breed west.
- Canvasback, *Aythya valisineria*. Locally com. WV on lakes and bays; uncom. SR on lakes east of Cascades.
- Redhead, *Aythya americana*. Com. PR on lakes east of Cascades, uncom. M and rare WV on fresh water west.
- Ring-necked Duck, *Aythya collaris*. Com. WV on fresh water, uncom. SR on wooded lakes in Cascades and east.
- Tufted Duck, *Aythya fuligula*. Rare WV, mostly west.
- Greater Scaup, *Aythya marila*. Abund. WV on salt water, locally com. on fresh water. Breeds in boreal forest.
- Lesser Scaup, *Aythya affinis*. Com. WV on fresh water, less so on salt water; uncom. SR on lakes east of Cascades, few west.
- Harlequin Duck, *Histrionicus histrionicus*. Com. WV (some in summer) on rocky coast; uncom. SR on mountain streams and rivers.
- Surf Scoter, *Melanitta perspicillata*. Abund. WV (some in summer) on salt water, rare M elsewhere. Breeds in boreal forest.
- White-winged Scoter, *Melanitta deglandi*. Abund. WV (some in summer) on salt water, rare M elsewhere. Breeds in boreal forest.
- Black Scoter, *Melanitta americana*. Locally com. WV on salt water. Breeds in Arctic and Subarctic.
- Long-tailed Duck, *Clangula hyemalis*. Com. WV on salt water. Breeds in Arctic.
- Bufflehead, *Bucephala albeola*. Abund. WV on salt and fresh water; few breed on lakes, mostly northeast.
- Common Goldeneye, *Bucephala clangula*. Com. WV on salt and fresh water; few breeding records on interior lakes.
- Barrow's Goldeneye, *Bucephala islandica*. Locally com. WV on salt water, uncom. M on fresh water; uncom. SR on mountain lakes.
- Hooded Merganser, *Lophodytes cucullatus*. Uncom. PR on fresh water, breeding on wooded lakes; some move to salt water in winter.
- Common Merganser, *Mergus merganser*. Com. WV on lakes, locally on salt water; uncom. SR on rivers.
- Red-breasted Merganser, *Mergus serrator*. Com. WV on salt water. Breeds in boreal forest and Arctic.
- Ruddy Duck, *Oxyura jamaicensis*. Com. WV on large lakes and bays; com. SR on lakes east of Cascades, few west.

Order GALLIFORMES

ODONTOPHORIDAE (New World quails)

Mountain Quail, *Oreortyx pictus*. Uncom. and local PR in wooded country around southern Puget Sound, lower Columbia River, and Blue Mountains.

Northern Bobwhite, *Colinus virginianus*. Uncom. and local PR, mostly Pierce and Thurston counties (introduced).

California Quail, *Callipepla californica*. Com. PR in lightly wooded and open country (introduced).

PHASIANIDAE (partridges, grouses, turkeys)

Wild Turkey, *Meleagris gallopavo*. Locally com. PR in wooded areas, increasing over much of east (introduced).

Ruffed Grouse, *Bonasa umbellus*. Uncom. PR in broadleaf forest.

Spruce Grouse, *Canachites canadensis*. Uncom. PR in conifer forest from Cascades east across northern part of state.

White-tailed Ptarmigan, *Lagopus leucura*. Uncom. PR in alpine zone of Cascades. Moves below tree line in winter.

Greater Sage-Grouse, *Centrocercus urophasianus*. Uncom., local and decreasing PR in sagebrush east of Cascades.

Dusky Grouse, *Dendragapus obscurus*. Uncom. PR in conifer forest east of Cascades.

Sooty Grouse, *Dendragapus fuliginosus*. Uncom. PR in conifer forest from Cascades west.

Sharp-tailed Grouse, *Tympanuchus phasianellus*. Uncom., local and decreasing PR in grassland east of Cascades, mostly northern part.

Gray Partridge, *Perdix perdix*. Uncom. PR in open country east of Cascades (introduced).

Ring-necked Pheasant, *Phasianus colchicus*. Com. PR in open country, less common west (introduced).

Chukar, *Alectoris chukar*. Uncom. PR in rocky open country east of Cascades (introduced).

Order PODICIPEDIFORMES

PODICIPEDIDAE (grebes)

Pied-billed Grebe, *Podilymbus podiceps*. Com. PR on lakes and ponds; uncom. WV on salt water.

Horned Grebe, *Podiceps auritus*. Com. WV on salt water and lakes; uncom. SR on lakes in northeast part.

Red-necked Grebe, *Podiceps grisegena*. Com. WV on salt water; locally com. SR on lakes in northeast part.

Eared Grebe, *Podiceps nigricollis*. Uncom. and local WV on salt water and lakes; locally com. SR in loose colonies on lakes east of Cascades.

Western Grebe, *Aechmophorus occidentalis*. Com. WV on large lakes and salt water; com. SR in loose colonies on few large lakes east of Cascades.

Clark's Grebe, *Aechmophorus clarkii*. Com. SR in loose colonies on few Columbia Basin lakes; rare WV on salt water.

Order COLUMBIFORMES

COLUMBIDAE (pigeons)

Rock Pigeon, *Columba livia*. Com. PR, with both domestic and feral populations, latter in cliffs of Columbia Basin (introduced).

Band-tailed Pigeon, *Patagioenas fasciata*. Com. SR and uncom. WV in forest west.

Eurasian Collared-Dove, *Streptopelia decaocto*. Locally com. PR; invaded from Southeast, has spread through much of state (introduced).

Mourning Dove, *Zenaida macroura*. Com. SR, occas. WV, in woodland and open country east, local and uncom. PR west.

Order CAPRIMULGIFORMES

CAPRIMULGIDAE (nightjars)

Common Nighthawk, *Chordeiles minor*. Com. SR (now uncom. west) of woodland and open country.

Common Poorwill, *Phalaenoptilus nuttallii*. Locally com. SR in dry conifer woodland east.

Order APODIFORMES

APODIDAE (swifts)

Black Swift, *Cypseloides niger*. Local and uncom. SR from Cascades west; breeds colonially on cliffs but ranges widely.

Vaux's Swift, *Chaetura vauxi*. Com. SR in forest, locally in cities.

White-throated Swift, *Aeronautes saxatilis*. Locally com. SR on cliffs of Columbia Basin.

TROCHILIDAE (hummingbirds)

Black-chinned Hummingbird, *Archilochus alexandri*. Uncom. SR in riparian woodland east of Cascades.

Anna's Hummingbird, *Calypte anna*. Locally com. PR west of Cascades, mostly in cities; slowly increasing east.

Calliope Hummingbird, *Selasphorus calliope*. Com. SR in dry conifer forest east of Cascades crest; fall M higher in mountains, rare M west.

Rufous Hummingbird, *Selasphorus rufus*. Com. SR in forest, fall M higher in mountains.

Order GRUIFORMES

RALLIDAE (rails)

Virginia Rail, *Rallus limicola*. Com. PR in freshwater marshes.

Sora, *Porzana carolina*. Com. SR, rare WV, in freshwater marshes.

American Coot, *Fulica americana*. Abund. WV on lakes and bays; com. SR in marshes, mostly east.

GRUIDAE (cranes)

Sandhill Crane, *Antigone canadensis*. Locally com. M east of Cascades and at Cape Flattery, from Arctic-breeding populations; wintering flock on lower Columbia River; few pairs breed in Klickitat and Yakima counties.

Order CHARADRIIFORMES

RECURVIROSTRIDAE (avocets & stilts)

Black-necked Stilt, *Himantopus mexicanus*. Locally com. SR in marshes east, rare M west.

American Avocet, *Recurvirostra americana*. Locally com. SR in marshes east, rare M west.

HAEMATOPODIDAE (oystercatchers)

Black Oystercatcher, *Haematopus bachmani*. Uncom. PR on rocky coasts.

CHARADRIIDAE (plovers)

Black-bellied Plover, *Pluvialis squatarola*. Com. WV and M on coast, uncom. fall M east of Cascades. Breeds in Arctic.

American Golden-Plover, *Pluvialis dominica*. Rare spring and uncom. fall M on coast; rare fall M east. Breeds in Arctic.

Pacific Golden-Plover, *Pluvialis fulva*. Rare spring and uncom. fall M on coast. Breeds in Alaska.

Snowy Plover, *Charadrius nivosus*. Uncom. SR on coast from Grays Harbor south.

Semipalmated Plover, *Charadrius semipalmatus*. Com. M, uncom. WV on coast; uncom. M east; has bred at Grays Harbor.

Killdeer, *Charadrius vociferus*. Com. PR west, com. SR and uncom. WV east.

SCOLOPACIDAE (sandpipers)

Whimbrel, *Numenius phaeopus*. Com. M, rare WV, on salt water; rare M at lakes east. Breeds in Arctic.

Long-billed Curlew, *Numenius americanus*. Uncom. and declining SR in open country east; rare M on salt water, one small wintering flock.

Bar-tailed Godwit, *Limosa lapponica*. Rare M, mostly fall, on coast. Breeds in Alaskan Arctic.

Hudsonian Godwit, *Limosa haemastica*. Rare M on coast. Breeds in Arctic.

Marbled Godwit, *Limosa fedoa*. Locally com. M and WV on coast, rare M east. Breeds in southern Alaska and on Great Plains.

Ruddy Turnstone, *Arenaria interpres*. Com. spring and uncom. fall M on coast. Breeds in Arctic.

Black Turnstone, *Arenaria melanocephala*. Com. WV on rocky coast. Breeds in Alaskan Arctic.

Red Knot, *Calidris canutus*. Locally com. spring and uncom. fall M on coast. Breeds in Arctic.

Surfbird, *Calidris virgata*. Com. WV on rocky coast. Breeds in Alaska mountains.

Ruff, *Calidris pugnax*. Rare fall M, mostly coastal mudflats. Breeds in Siberian Arctic.

Sharp-tailed Sandpiper, *Calidris acuminata*. Rare fall M on coast. Breeds in Siberian Arctic.

Stilt Sandpiper, *Calidris himantopus*. Rare fall M on fresh and salt water. Breeds in Arctic.

Sanderling, *Calidris alba*. Com. M and WV on coast; uncom. M, mostly fall, east. Breeds in Arctic.

Dunlin, *Calidris alpina*. Abund. WV and M on coast; uncom. M at lakes east. Breeds in Arctic.

Rock Sandpiper, *Calidris ptilocnemis*. Uncom. and declining WV on rocky coast. Breeds in Alaskan Arctic.

Baird's Sandpiper, *Calidris bairdii*. Uncom. M, mostly fall, on fresh and salt water. Breeds in Arctic.

Least Sandpiper, *Calidris minutilla*. Com. M and uncom. WV on coast, com. M at lakes and marshes east. Breeds in boreal forest.

- Buff-breasted Sandpiper, *Calidris subruficollis*. Rare fall M, mostly coastal dunes. Breeds in Arctic.
- Pectoral Sandpiper, *Calidris melanotos*. Uncom. M, mostly fall, on fresh and salt water. Breeds in Arctic.
- Semipalmated Sandpiper, *Calidris pusilla*. Uncom. M, mostly on fresh water and more common east. Breeds in Arctic.
- Western Sandpiper, *Calidris mauri*. Abund. M and uncom. WV on coast, com. M at lakes east. Breeds in Alaskan Arctic.
- Short-billed Dowitcher, *Limnodromus griseus*. Abund. M on coast, rare at freshwater elsewhere. Breeds in boreal forest.
- Long-billed Dowitcher, *Limnodromus scolopaceus*. Com. M and uncom. WV on fresh and salt water. Breeds in Alaskan Arctic.
- Wilson's Snipe, *Gallinago delicata*. Com. PR west, SR east, in marshes.
- Spotted Sandpiper, *Actitis macularius*. Com. SR on rivers and lakes; uncom. WV on coastal fresh and salt water.
- Solitary Sandpiper, *Tringa solitaria*. Rare M on fresh water. Breeds in boreal forest.
- Wandering Tattler, *Tringa incana*. Uncom. M on rocky coasts. Breeds in Alaska.
- Lesser Yellowlegs, *Tringa flavipes*. Uncom. spring and com. fall M, mostly fresh water. Breeds in boreal forest and Subarctic.
- Willet, *Tringa semipalmata*. Rare and local WV on coast. Breeds to east and south.
- Greater Yellowlegs, *Tringa melanoleuca*. Com. M on fresh and salt water, local WV on coast. Breeds in boreal forest.
- Wilson's Phalarope, *Phalaropus tricolor*. Com. SR in marshes east, rare M on coast.
- Red-necked Phalarope, *Phalaropus lobatus*. Com. M offshore and in protected salt water, less common in fresh water. Breeds in Arctic.
- Red Phalarope, *Phalaropus fulicarius*. Com. M offshore, rarely to protected waters and rarely winters. Breeds in Arctic.
- STERCORARIIDAE (skuas)
- South Polar Skua, *Stercorarius maccormicki*. Rare SV offshore. Breeds in Antarctica.
- Pomarine Jaeger, *Stercorarius pomarinus*. Uncom. M offshore. Breeds in Arctic.
- Parasitic Jaeger, *Stercorarius parasiticus*. Com. M offshore and along coast. Breeds in Arctic.
- Long-tailed Jaeger, *Stercorarius longicaudus*. Uncom. fall M offshore. Breeds in Arctic.
- ALCIDAE (alcids)
- Common Murre, *Uria aalge*. Com. WV on coast; declining SR on outer-coastal islands; colonial breeder.
- Pigeon Guillemot, *Cephus columba*. Com. PR on protected salt water.
- Marbled Murrelet, *Brachyramphus marmoratus*. Uncom. and declining PR on coast; breeds in old-growth conifer forest.
- Scripps's Murrelet, *Synthliboramphus scrippsi*. Rare SV offshore.
- Ancient Murrelet, *Synthliboramphus antiquus*. Com. WV (Nov.-Jan. in protected waters) on coast; has bred on offshore islands.
- Cassin's Auklet, *Ptychoramphus aleuticus*. Com. but declining PR on outer coast; colonial breeder.
- Parakeet Auklet, *Aethia psittacula*. Rare WV offshore.
- Rhinoceros Auklet, *Cerorhinca monocerata*. Abund. SR and locally common WV on coast; colonial breeder.
- Horned Puffin, *Fratercula corniculata*. Rare WV, mostly offshore.
- Tufted Puffin, *Fratercula cirrhata*. Com. SR on coast, winters offshore; colonial breeder.
- LARIDAE (gulls & terns)
- Black-legged Kittiwake, *Rissa tridactyla*. Uncom. WV (some summer) on outer coast. Breeds around Bering Sea.
- Sabine's Gull, *Xema sabini*. Com. M offshore. Breeds in Arctic.
- Bonaparte's Gull, *Chroicocephalus philadelphia*. Com. M and uncom. WV on salt water, rare M on lakes east. Breeds in boreal forest.
- Franklin's Gull, *Leucophaeus pipixcan*. Uncom. fall M on fresh and salt water west; rare spring M on lakes east. Breeds to east and south.
- Heermann's Gull, *Larus heermanni*. Com. SV and fall M, rarely WV, on coast. Breeds in Gulf of California.
- Short-billed Gull, *Larus brachyrhynchus*. Com. WV on coast, much less so inland. Breeds in boreal forest.

- Ring-billed Gull, *Larus delawarensis*. Com. SR at Grays Harbor and on lakes east, colonial breeder; com. WV on fresh water, much less so on salt water.
- Western Gull, *Larus occidentalis*. Com. PR on outer coast, decreasing east through protected waters; colonial breeder.
- California Gull, *Larus californicus*. Abund. fall M, much less common in spring and winter, on coast; com. SR on lakes east, colonial breeder.
- Herring Gull, *Larus argentatus*. Uncom. WV on fresh and salt water. Breeds in boreal forest.
- Iceland Gull, *Larus glaucoides*. Locally com. WV on salt and coastal fresh water. Breeds in Arctic.
- Lesser Black-backed Gull, *Larus fuscus*. Rare WV.
- Glaucous-winged Gull, *Larus glaucescens*. Com. PR on coast, colonial breeder; rare breeder and WV in interior.
- Glaucous Gull, *Larus hyperboreus*. Rare WV on fresh and salt water. Breeds in Arctic.
- Caspian Tern, *Hydroprogne caspia*. Locally com. SR on coast, Columbia River, and interior reservoirs; colonial breeder.
- Black Tern, *Chlidonias niger*. Locally com. SR on lakes east; rare M on fresh and salt water west.
- Common Tern, *Sterna hirundo*. Com. but declining M on coast, rare M on lakes east. Breeds to northeast in boreal forest.
- Arctic Tern, *Sterna paradisaea*. Com. M offshore, occasionally on coast. Breeds in Arctic (few pairs at Dungeness).
- Forster's Tern, *Sterna forsteri*. Locally com. SR on lakes east; rare M on fresh and salt water west.
- Elegant Tern, *Thalasseus elegans*. Irregular fall M on coast (usually absent but has been com. during some El Niño years, few recent records). Breeds in Gulf of California.

Order GAVIIFORMES

GAVIIDAE (loons)

- Red-throated Loon, *Gavia stellata*. Com. WV on salt water. Breeds in Arctic and Subarctic.
- Pacific Loon, *Gavia pacifica*. Com. WV on salt water; abund. M on outer coast. Breeds in Arctic.
- Common Loon, *Gavia immer*. Com. WV on salt and fresh water; rare breeder on undisturbed lakes.
- Yellow-billed Loon, *Gavia adamsii*. Rare WV on salt water. Breeds in Arctic.

Order PROCELLARIIFORMES

DIOMEDEIDAE (albatrosses)

- Laysan Albatross, *Phoebastria immutabilis*. Rare offshore, mostly WV. Breeds in Hawaiian islands.
- Black-footed Albatross, *Phoebastria nigripes*. Com. SV and rare WV offshore. Breeds in Hawaiian Islands.
- Short-tailed Albatross, *Phoebastria albatrus*. Rare offshore visitor. Breeds sparingly on W Pacific islands.

HYDROBATIDAE (storm-petrels)

- Fork-tailed Storm-Petrel, *Hydrobates furcatus*. Com. SR on outer coast, breeding on islands and foraging offshore.
- Leach's Storm-Petrel, *Hydrobates leucorhous*. Com. SR on outer coast, breeding on islands and foraging well offshore.

PROCELLARIIDAE (petrels)

- Northern Fulmar, *Fulmarus glacialis*. Com. WV and uncom. SV offshore. Breeds around Bering Sea.
- Mottled Petrel, *Pterodroma inexpectata*. Rare offshore, mostly WV. Breeds in New Zealand.
- Buller's Shearwater, *Ardenna bulleri*. Uncom. SV (autumn) offshore. Breeds in New Zealand.
- Short-tailed Shearwater, *Ardenna tenuirostris*. Uncom. M, mostly fall, offshore, rarely into Puget Sound. Breeds in Australia.
- Sooty Shearwater, *Ardenna grisea*. Com. SV on outer coast, rarely into Puget Sound. Breeds in New Zealand.
- Pink-footed Shearwater, *Ardenna creatopus*. Com. SV offshore. Breeds in Chile.
- Flesh-footed Shearwater, *Ardenna carneipes*. Rare SV offshore. Breeds in Australia.

Manx Shearwater, *Puffinus puffinus*. Rare SV on outer coast. Breeds in North Atlantic, possibly North Pacific.

Order SULIFORMES

SULIDAE (boobies)

Brown Booby, *Sula leucogaster*. Rare visitor from tropical ocean.

PHALACROCORACIDAE (cormorants)

Brandt's Cormorant, *Urile penicillatus*. Com. WV and abund. M (from south) on coast, immatures remain through summer; colonial breeder on rocks in Pacific County, occasionally elsewhere.

Pelagic Cormorant, *Urile pelagicus*. Com. PR on coast; colonial breeder.

Double-crested Cormorant, *Nannopterum auritum*. Com. PR on salt water and large interior lakes and rivers; colonial breeder.

Order PELECANIFORMES

PELECANIDAE (pelicans)

American White Pelican, *Pelecanus erythrorhynchos*. Uncom. SV on lakes of Columbia Basin, local colonial breeder and rare WV on Columbia River; regular visitor west at present.

Brown Pelican, *Pelecanus occidentalis*. Com. SV on outer coast, rarely into protected waters. Breeds from California south.

ARDEIDAE (herons)

American Bittern, *Botaurus lentiginosus*. Uncom. PR in freshwater marshes.

Great Blue Heron, *Ardea herodias*. Com. PR on fresh and salt water; colonial breeder.

Great Egret, *Ardea alba*. Growing breeding colonies in Grant County, probably elsewhere; otherwise uncommon but increasing SV (and rarer WV) on fresh and salt water, mostly on coast and in Columbia Basin.

Green Heron, *Butorides virescens*. Uncom. SR (rarely winters) on streams and swamps west of Cascades.

Black-crowned Night-Heron, *Nycticorax nycticorax*. Locally com. SR in marshes of Columbia Basin; colonial breeder; rare and local WV east and west.

THRESKIORNITHIDAE (ibises and spoonbills)

White-faced Ibis, *Plegadis chihi*. Rare SV, mostly in Columbia Basin; has attempted to breed.

Order CATHARTIFORMES

CATHARTIDAE (American vultures)

Turkey Vulture, *Cathartes aura*. Uncom. SR in most parts of state; large flocks at times seen in coastal migration.

Order ACCIPITRIFORMES

PANDIONIDAE (osprey)

Osprey, *Pandion haliaetus*. Uncom. SR on fresh and salt water.

ACCIPITRIDAE (hawks)

White-tailed Kite, *Elanus leucurus*. Rare PR at woodland edge southwest.

Golden Eagle, *Aquila chrysaetos*. Uncom. PR in mountains and dry canyons.

Northern Harrier, *Circus hudsonius*. Uncom. PR in open country.

Sharp-shinned Hawk, *Accipiter striatus*. Uncom. SR in mountain forest, more often seen WV in wooded lowlands.

Cooper's Hawk, *Accipiter cooperii*. Uncom. PR in mixed and broadleaf forests.

Northern Goshawk, *Accipiter gentilis*. Rare PR in mature conifer forest.

Bald Eagle, *Haliaeetus leucocephalus*. Com. PR on fresh and salt water.

Red-shouldered Hawk, *Buteo lineatus*. Rare PR in woodland, mostly southwest. No confirmed breeding record.
 Broad-winged Hawk, *Buteo platypterus*. Rare M, spring on outer coast, fall in Cascades.
 Swainson's Hawk, *Buteo swainsoni*. Uncom. SR in open country east of Cascades.
 Red-tailed Hawk, *Buteo jamaicensis*. Com. PR in all habitats, numbers augmented in winter.
 Rough-legged Hawk, *Buteo lagopus*. Uncom. to locally com. WV in open country. Breeds in Arctic.
 Ferruginous Hawk, *Buteo regalis*. Rare and local SR in southeast quarter of state, mostly in canyons.

Order STRIGIFORMES

TYTONIDAE (barn owls)

Barn Owl, *Tyto alba*. Uncom. PR in any habitat that provides nest sites.

STRIGIDAE (typical owls)

Flammulated Owl, *Psilosops flammeolus*. Uncom. SR in dry conifer forest east.

Western Screech-Owl, *Megascops kennicottii*. Uncom. PR in forest.

Great Horned Owl, *Bubo virginianus*. Com. PR in all habitats.

Snowy Owl, *Bubo scandiacus*. Uncom. to rare (erratic) WV in open country. Breeds in Arctic.

Northern Hawk Owl, *Surnia ulula*. Rare WV, usually associated with open woodland; has bred in Okanogan County.

Northern Pygmy-Owl, *Glaucidium gnoma*. Uncom. PR in conifer forest.

Burrowing Owl, *Athene cunicularia*. Uncom. and declining SR in sagebrush steppe east.

Spotted Owl, *Strix occidentalis*. Uncom. and declining PR of old-growth conifer forest west.

Barred Owl, *Strix varia*. Uncom. but increasing PR in forest.

Great Gray Owl, *Strix nebulosa*. Rare and local PR in open conifer forest; breeds east, more widespread in winter.

Long-eared Owl, *Asio otus*. Uncom. PR (more local in winter) in forest, especially riparian woodland; much less common west.

Short-eared Owl, *Asio flammeus*. Uncom. PR in open country.

Boreal Owl, *Aegolius funereus*. Rare PR in high-elevation conifer forest.

Northern Saw-whet Owl, *Aegolius acadicus*. Com. but rarely seen PR in forest.

Order CORACIIFORMES

ALCEDINIDAE (kingfishers)

Belted Kingfisher, *Megaceryle alcyon*. Uncom. PR on fresh and salt water, more widespread in winter.

Order PICIFORMES

PICIDAE (woodpeckers)

Lewis's Woodpecker, *Melanerpes lewis*. Uncom. SR (winters locally) in ponderosa pineland and oak and riparian woodland east of Cascades; rare M west.

Acorn Woodpecker, *Melanerpes formicivorus*. Rare PR in oak woodland in southern Klickitat County.

Williamson's Sapsucker, *Sphyrapicus thyroideus*. Uncom. SR in mature dry conifer forest east of Cascades crest; rare M west.

Red-naped Sapsucker, *Sphyrapicus nuchalis*. Com. SR in forest east of Cascades crest; rare M west.

Red-breasted Sapsucker, *Sphyrapicus ruber*. Uncom. PR in mature forest from Cascades west, local on east side of Cascades.

American Three-toed Woodpecker, *Picoides dorsalis*. Uncom. PR in high-elevation conifer forest east of Cascades crest.

Black-backed Woodpecker, *Picoides arcticus*. Uncom. PR in conifer forest east of Cascades crest; mostly in burns.

Downy Woodpecker, *Dryobates pubescens*. Com. PR in broadleaf and mixed forest.

Hairy Woodpecker, *Dryobates villosus*. Com. PR in conifer and mixed forest.

White-headed Woodpecker, *Dryobates albolarvatus*. Uncom. PR in ponderosa pineland from Cascades east.
 Northern Flicker, *Colaptes auratus*. Com. PR in all wooded habitats.
 Pileated Woodpecker, *Dryocopus pileatus*. Uncom. PR in mature forest, increasingly in suburbs.

Order FALCONIFORMES

FALCONIDAE (falcons)

American Kestrel, *Falco sparverius*. Com. SR and uncom. WV in open country and woodland east of Cascades, much less common west; also SR above treeline in mountains.
 Merlin, *Falco columbarius*. Uncom. WV, especially coast, mostly M east; increasingly breeding in forests and cities west.
 Gyrfalcon, *Falco rusticolus*. Rare WV in open country. Breeds in Arctic.
 Peregrine Falcon, *Falco peregrinus*. Uncom. PR where nesting cliffs or tall humanmade structures available in Cascades and west; uncom. WV elsewhere, especially coastal areas; slightly more common in migration.
 Prairie Falcon, *Falco mexicanus*. Uncom. PR (less common in winter) in canyons east of Cascades; rarely winters west.

Order PASSERIFORMES

TYRANNIDAE (tyrant flycatchers)

Ash-throated Flycatcher, *Myiarchus cinerascens*. Uncom. SR in riparian woodland in Kittitas, Yakima and Klickitat counties.
 Tropical Kingbird, *Tyrannus melancholicus*. Rare late fall M, mostly to coastal areas. Breeds in American tropics.
 Western Kingbird, *Tyrannus verticalis*. Com. SR in open country east of Cascades; very local SR west.
 Eastern Kingbird, *Tyrannus tyrannus*. Com. SR in open country, usually near water, east of Cascades; very local SR west.
 Olive-sided Flycatcher, *Contopus cooperi*. Uncom. SR in forest.
 Western Wood-Pewee, *Contopus sordidulus*. Com. SR (less common west) in forest and woodland.
 Willow Flycatcher, *Empidonax traillii*. Com. SR in broadleaf forest and shrubland, usually near water.
 Least Flycatcher, *Empidonax minimus*. Rare SR in broadleaf forest, mostly northeast.
 Hammond's Flycatcher, *Empidonax hammondi*. Com. SR in mature forest.
 Gray Flycatcher, *Empidonax wrightii*. Com. SR in open ponderosa pineland east of Cascades crest.
 Dusky Flycatcher, *Empidonax oberholseri*. Com. SR in dry conifer forest east of Cascades crest, rare M west.
 Pacific-slope Flycatcher, *Empidonax difficilis*. Com. SR in mature forest.
 Black Phoebe, *Sayornis nigricans*. Rare but increasing PR west; breeding locally.
 Say's Phoebe, *Sayornis saya*. Com. SR in open country east of Cascades; rare M west.

VIREONIDAE (vireos)

Hutton's Vireo, *Vireo huttoni*. Uncom. PR in broadleaf and mixed forest, mostly near coast.
 Cassin's Vireo, *Vireo cassinii*. Com. SR in conifer forest.
 Warbling Vireo, *Vireo gilvus*. Com. SR in low broadleaf forest.
 Red-eyed Vireo, *Vireo olivaceus*. Com. SR in mature riparian woodland.

LANIIDAE (shrikes)

Loggerhead Shrike, *Lanius ludovicianus*. Uncom. SR and rare WV in sagebrush steppe east of Cascades; rare M west.
 Northern Shrike, *Lanius borealis*. Uncom. WV in open country, more common east. Breeds in boreal forest.

CORVIDAE (crows & jays)

Canada Jay, *Perisoreus canadensis*. Com. PR in conifer forest, mostly in mountains.
 Steller's Jay, *Cyanocitta stelleri*. Com. PR in conifer forest.
 Blue Jay, *Cyanocitta cristata*. Rare WV, usually at bird feeders.

- California Scrub-Jay, *Aphelocoma californica*. Com. PR in oak woodland and urban/suburban areas southwest, east to Klickitat County; slowly expanding range northward in Puget Sound lowlands, breeds north to King County.
- Clark's Nutcracker, *Nucifraga columbiana*. Com. PR in alpine zone east of Cascades crest, less common in ponderosa pineland.
- Black-billed Magpie, *Pica hudsonia*. Com. PR in riparian woodland and open country east of Cascades.
- American Crow, *Corvus brachyrhynchos*. Com. PR in all wooded habitats.
- Common Raven, *Corvus corax*. Com. PR on coast, in mountains and in open country east of Cascades.
- PARIDAE (tits)
- Black-capped Chickadee, *Poecile atricapillus*. Com. PR in broadleaf forest.
- Mountain Chickadee, *Poecile gambeli*. Com. PR in conifer forest east of Cascades crest.
- Chestnut-backed Chickadee, *Poecile rufescens*. Com. PR in wet conifer forest in mountains and western lowlands.
- Boreal Chickadee, *Poecile hudsonicus*. Uncom. PR in high-elevation conifer forest along northern border east of Cascades crest.
- ALAUDIDAE (larks)
- Horned Lark, *Eremophila alpestris*. Com. SR and abund. WV in open country east of Cascades; uncom. and local PR on prairies and coastal dunes west; uncom. SV in alpine zone.
- HIRUNDINIDAE (swallows)
- Purple Martin, *Progne subis*. Uncom. SR west, mostly around Puget Sound.
- Tree Swallow, *Tachycineta bicolor*. Com. SR in wooded habitats near water.
- Violet-green Swallow, *Tachycineta thalassina*. Com. SR in forest and canyon country, locally in cities.
- Northern Rough-winged Swallow, *Stelgidopteryx serripennis*. Com. SR, usually near fresh or salt water where earth banks available for nesting.
- Bank Swallow, *Riparia riparia*. Locally com. SR east of Cascades where sand banks available for colonial nesting; very local breeder and rare M west.
- Cliff Swallow, *Petrochelidon pyrrhonota*. Com. SR where cliffs, bridges or buildings available for colonial nesting.
- Barn Swallow, *Hirundo rustica*. Com. SR where bridges or buildings available for nesting.
- AEGITHALIDAE (long-tailed tits)
- Bushtit, *Psaltriparus minimus*. Com. PR in broadleaf forest and shrubland west of Cascades, also local in riparian woodland in Kittitas and Yakima counties; colonizing Columbia Basin.
- REGULIDAE (kinglets)
- Ruby-crowned Kinglet, *Corthylio calendula*. Com. WV in wooded habitats of lowlands; com. SR in broadleaf thickets in conifer forest east of Cascades crest.
- Golden-crowned Kinglet, *Regulus satrapa*. Com. PR in conifer forest, more widespread in winter.
- BOMBYCILLIDAE (waxwings)
- Bohemian Waxwing, *Bombycilla garrulus*. Com. WV east of Cascades, mostly in cities; irregular and uncom. WV west; has bred in conifer forest along northern border.
- Cedar Waxwing, *Bombycilla cedrorum*. Com. SR, much less com. WV (more common east), in wooded habitats.
- SITTIDAE (nuthatches)
- Red-breasted Nuthatch, *Sitta canadensis*. Com. PR in conifer forest, more widespread in winter.
- White-breasted Nuthatch, *Sitta carolinensis*. Com. PR in riparian woodland and ponderosa pineland east of Cascades crest; uncom. PR in oak woodland south of Puget Sound.
- Pygmy Nuthatch, *Sitta pygmaea*. Locally com. PR in ponderosa pineland east of Cascades crest.
- CERTHIIDAE (creepers)
- Brown Creeper, *Certhia americana*. Com. PR in conifer forest, more widespread in winter.
- TROGLODYTIDAE (wrens)
- Rock Wren, *Salpinctes obsoletus*. Com. SR in rocky areas, mostly in open country, east of Cascades.
- Canyon Wren, *Catherpes mexicanus*. Uncom. PR in rocky canyons, both open and wooded, east of Cascades.

House Wren, *Troglodytes aedon*. Com. SR in riparian woodland east of Cascades; locally com. SR, mostly in oak woodland, west.

Pacific Wren, *Troglodytes pacificus*. Com. PR in conifer forest, more widespread in winter.

Marsh Wren, *Cistothorus palustris*. Com. PR (east of Cascades, most migrate south) in freshwater marshes.

Bewick's Wren, *Thryomanes bewickii*. Com. PR in broadleaf forest and shrubland west of Cascades, rapidly expanding up Columbia River drainage in east.

MIMIDAE (mockingbirds)

Gray Catbird, *Dumetella carolinensis*. Com. SR in riparian woodland east of Cascades.

Sage Thrasher, *Oreoscoptes montanus*. Com. SR in sagebrush steppe east of Cascades; rare M west.

Northern Mockingbird, *Mimus polyglottos*. Rare PR, mostly vagrant individuals in woodland or urban areas, but has bred a few times east.

STURNIDAE (starlings)

European Starling, *Sturnus vulgaris*. Com. PR in all habitats that provide nesting holes (introduced).

CINCLIDAE (dippers)

American Dipper, *Cinclus mexicanus*. Uncom. PR on wooded rivers and streams.

TURDIDAE (thrushes)

Western Bluebird, *Sialia mexicana*. Locally com. SR in open woodland east of Cascades crest; even more local, mostly in oak woodland, west.

Mountain Bluebird, *Sialia currucoides*. Com. SR in alpine zone, open woodland and open country east of Cascades crest; rare M west.

Townsend's Solitaire, *Myadestes townsendi*. Com. SR in mountain conifer forest, uncom. M elsewhere; uncom. WV in lowlands, associated with junipers; widespread uncom. M west.

Veery, *Catharus fuscescens*. Com. SR in broadleaf forest east of Cascades crest.

Swainson's Thrush, *Catharus ustulatus*. Com. SR in forest (only in wet conifer forest east of Cascades crest).

Hermit Thrush, *Catharus guttatus*. Com. SR in mountain conifer forest, uncom. M elsewhere; uncom. WV in forest near coast.

American Robin, *Turdus migratorius*. Com. SR in all habitats with trees and shrubs; withdraws from mountains and becomes abundant in lowlands in winter.

Varied Thrush, *Ixoreus naevius*. Com. SR in mountain conifer forest; com. WV in lowland forest, local east.

PASSERIDAE (sparrows)

House Sparrow, *Passer domesticus*. Com. PR in settled areas, more in farmland east (introduced).

MOTACILLIDAE (wagtails)

American Pipit, *Anthus rubescens*. Com. M in open country; com. SR in wet meadows in mountains; uncom. WV in western lowlands.

FRINGILLIDAE (finches)

Evening Grosbeak, *Coccothraustes vespertinus*. Com. but declining PR in forest, irregular in movements.

Pine Grosbeak, *Pinicola enucleator*. Uncom. PR in mountain conifer forest, wandering to lowlands in winter.

Gray-crowned Rosy-Finch, *Leucosticte tephrocotis*. Uncom. SR in alpine zone in mountains; locally com. WV in open country with canyons east.

House Finch, *Haemorhous mexicanus*. Com. PR in riparian woodland and settled areas, also nearby open country in winter.

Purple Finch, *Haemorhous purpureus*. Com. PR in forest from east slope of Cascades west.

Cassin's Finch, *Haemorhous cassinii*. Com. SR (few remain in winter) in conifer forest east of Cascades crest.

Common Redpoll, *Acanthis flammea*. Uncom. WV in open country east, rarely west. Breeds in Arctic and Subarctic.

Red Crossbill, *Loxia curvirostra*. Com. PR in conifer forest; numbers fluctuate from year to year.

White-winged Crossbill, *Loxia leucoptera*. Irregular PR in mountain spruce forest, varying from absent to common; no evidence of breeding.

Pine Siskin, *Spinus pinus*. Com. PR in conifer and mixed forest, more widespread in winter.

Lesser Goldfinch, *Spinus psaltria*. Uncom. PR in oak woodland in Klickitat County, extending northward both west and east.

American Goldfinch, *Spinus tristis*. Com. SR in broadleaf and riparian woodland, widespread but less common in winter.

CALCARIIDAE (longspurs and snow buntings)

Lapland Longspur, *Calcarius lapponicus*. Uncom. M and WV in open country. Breeds in Arctic.

Snow Bunting, *Plectrophenax nivalis*. Uncom. and local WV in open country, more common east. Breeds in Arctic.

PASSERELLIDAE (American sparrows)

Green-tailed Towhee, *Pipilo chlorurus*. Rare SR in shrubs in open conifer forest on south slope of Blue Mountains.

Spotted Towhee, *Pipilo maculatus*. Com. PR west and SR east in shrublands.

American Tree Sparrow, *Spizelloides arborea*. Uncom. WV in open country east, less common west. Breeds in Arctic-boreal forest transition.

Chipping Sparrow, *Spizella passerina*. Com. SR in open conifer forest east of Cascades crest, locally com. SR in open woodland in San Juan County and south of Puget Sound west.

Clay-colored Sparrow, *Spizella pallida*. Rare SR in open woodland in northeast corner; has bred in Okanogan and Spokane counties; rare WV west.

Brewer's Sparrow, *Spizella breweri*. Com. SR in sagebrush steppe east of Cascades.

Vesper Sparrow, *Pooecetes gramineus*. Com. SR in grassland and shrub steppe east of Cascades; local, uncom. and declining SR in prairies west.

Lark Sparrow, *Chondestes grammacus*. Com. SR in sagebrush steppe and uncom. SR in open conifer forest east of Cascades; rare M west.

Black-throated Sparrow, *Amphispiza bilineata*. Rare and very local SR in sagebrush steppe.

Sagebrush Sparrow, *Artemisospiza nevadensis*. Uncom. SR in sagebrush steppe of Columbia Basin; rare M west.

Savannah Sparrow, *Passerculus sandwichensis*. Com. SR and abund. M in moist grassland and farmland; uncom. WV in farmland west.

Grasshopper Sparrow, *Ammodramus savannarum*. Uncom. SR in dry grassland and open sagebrush steppe east of Cascades.

Fox Sparrow, *Passerella iliaca*. Com. WV in shrublands west of Cascades, much less com. east; locally com. SR in shrubby clearings in mountains east of Cascades crest and on tip of Olympic Peninsula.

Song Sparrow, *Melospiza melodia*. Com. PR in shrublands (usually associated with water east of Cascades).

Lincoln's Sparrow, *Melospiza lincolnii*. Com. M and uncom. WV in shrublands west of Cascades; locally com. SR in shrubby wetlands in mountains.

Swamp Sparrow, *Melospiza georgiana*. Rare WV in shrublands near water, mostly west.

White-throated Sparrow, *Zonotrichia albicollis*. Rare WV in forest and shrublands, mostly west.

Harris's Sparrow, *Zonotrichia querula*. Rare WV in shrublands.

White-crowned Sparrow, *Zonotrichia leucophrys*. Com. SR in shrublands from east side of Cascades west to coast (coastal subspecies); com. WV and abundant M in lowlands throughout, also breeding in North Cascades (subarctic subspecies); uncom. SR in high mountains at eastern edge of state (Rocky Mountain subspecies).

Golden-crowned Sparrow, *Zonotrichia coronata*. Com. M and locally com. WV in shrublands from coast to east side of Cascades, rarer still farther east; has bred in north Cascades.

Dark-eyed Junco, *Junco hyemalis*. Com. PR in conifer forest, widespread in most habitats in winter.

ICTERIIDAE (Yellow-breasted Chat)

Yellow-breasted Chat, *Icteria virens*. Uncom. SR in shrubby riparian woodland east of Cascades; has summered west.

ICTERIDAE (troupials)

Yellow-headed Blackbird, *Xanthocephalus xanthocephalus*. Com. SR in marshes east of Cascades, rare WV at feedlots; rare M, occas. SR, west.

Bobolink, *Dolichonyx oryzivorus*. Local and uncom. SR in hay meadows east of Cascades; rare M west.

Western Meadowlark, *Sturnella neglecta*. Com. SR (few winter) in open country east; uncom. and local PR in open country west, more in winter.

Bullock's Oriole, *Icterus bullockii*. Com. SR in riparian woodland, less common west.

Red-winged Blackbird, *Agelaius phoeniceus*. Com. SR in shrubby wetlands and marshes; locally com. WV in farmland.

Tricolored Blackbird, *Agelaius tricolor*. Very local SR in Columbia Basin marshes, rare WV southwest.

Brown-headed Cowbird, *Molothrus ater*. Com. SR in all habitats; local and uncom. WV in farmland.

Rusty Blackbird, *Euphagus carolinus*. Rare WV.

Brewer's Blackbird, *Euphagus cyanocephalus*. Com. SR in settled and open country east, much more local west; locally com. WV in farmland and cities.

PARULIDAE (wood warblers)

Northern Waterthrush, *Parkesia noveboracensis*. Uncom. SR in wooded wetlands from Okanogan County east; rare M elsewhere; occasional WV west.

Black-and-white Warbler, *Mniotilta varia*. Rare M.

Tennessee Warbler, *Leiothlypis peregrina*. Rare M, even rarer WV.

Orange-crowned Warbler, *Leiothlypis celata*. Com. SR in open broadleaf forest and shrubland west, much more local east in mountain forest.

Nashville Warbler, *Leiothlypis ruficapilla*. Com. SR in broadleaf thickets in dry conifer forest east of Cascades crest; rare M west.

MacGillivray's Warbler, *Geothlypis tolmiei*. Com. SR in riparian thickets east of Cascades, more local in shrubland, especially revegetating clearings, west.

Common Yellowthroat, *Geothlypis trichas*. Com. SR in shrubby wetlands (very local east).

American Redstart, *Setophaga ruticilla*. Uncom. SR in broadleaf forest east of Cascades (more common to north), very local west.

Yellow Warbler, *Setophaga petechia*. Com. SR in broadleaf and riparian woodland.

Chestnut-sided Warbler, *Setophaga pensylvanica*. Rare M.

Palm Warbler, *Setophaga palmarum*. Rare WV, mostly on outer coast.

Yellow-rumped Warbler, *Setophaga coronata*. Com. SR in mountain conifer forest; com. M and locally com. WV in lowland wooded habitats.

Black-throated Gray Warbler, *Setophaga nigrescens*. Com. SR in broadleaf and mixed forest west of Cascades; local on east slope.

Townsend's Warbler, *Setophaga townsendi*. Com. SR in mountain conifer forest; uncom. M and WV in forest near coast. Freely hybridizes with Hermit Warbler.

Hermit Warbler, *Setophaga occidentalis*. Com. SR in conifer forest in south Cascades, Willapa Hills and east side of Olympics. Freely hybridizes with Townsend's Warbler.

Wilson's Warbler, *Cardellina pusilla*. Com. SR in forest, more local east.

CARDINALIDAE (cardinals and allies)

Western Tanager, *Piranga ludoviciana*. Com. SR in conifer and mixed forest.

Rose-breasted Grosbeak, *Pheucticus ludovicianus*. Rare M, occasional WV.

Black-headed Grosbeak, *Pheucticus melanocephalus*. Com. SR in broadleaf forest.

Lazuli Bunting, *Passerina amoena*. Com. SR on broadleaf shrubby hillsides east of Cascades crest, very local west.

IRREGULAR SPECIES

These species, not recorded annually and thus very rarely seen in Washington, are included to show the great dispersal abilities of birds. Note they come from all directions, as can be seen from their breeding sources: A - arctic N. America; AK - Alaska; Ant - Antarctica; Atl - Atlantic Ocean; E - eastern/central N. America; EP - east Pacific Ocean; Eur - Eurasia (mostly from Siberia); GP - Great Plains; N - northern N. America; NA - north Atlantic Ocean; NC - north-central North America; NE - northeastern N. America; NGP - northern Great Plains; NP - north Pacific Ocean; NWP - northwest Pacific Ocean; RM - Rocky Mountains; SA - South America; SE - southeastern N. America; S - southern N. America; SGP - southern Great Plains; SP - south Pacific Ocean; SW - southwestern N. America; TP - tropical Pacific Ocean; WP - western Pacific Ocean.

Fulvous Whistling-Duck, *Dendrocygna bicolor* (SW)

Taiga Bean-Goose, *Anser fabalis* (Eur)
 Whooper Swan, *Cygnus cygnus* (Eur)
 Baikal Teal, *Sibirionetta formosa* (Eur)
 Garganey, *Spatula querquedula* (Eur)
 Falcated Duck, *Mareca falcata* (Eur)
 Steller's Eider, *Polysticta stelleri* (AK)
 King Eider, *Somateria spectabilis* (A)
 Common Eider, *Somateria mollissima* (A)
 Smew, *Mergellus albellus* (Eur)
 White-winged Dove, *Zenaida asiatica* (SW)
 Yellow-billed Cuckoo, *Coccyzus americanus* (SW/E)
 Black-billed Cuckoo, *Coccyzus erythrophthalmus* (NE)
 Costa's Hummingbird, *Calypte costae* (SW)
 Allen's Hummingbird, *Selasphorus sasin* (SW)
 Broad-tailed Hummingbird, *Selasphorus platycercus* (RM)
 Broad-billed Hummingbird, *Cynanthus latirostris* (SW)
 Purple Gallinule, *Porphyrio martinicus* (E)
 Yellow Rail, *Coturnicops noveboracensis* (N)
 Lesser Sand-Plover, *Charadrius mongolus* (Eur)
 Wilson's Plover, *Charadrius wilsonia* (SE)
 Common Ringed Plover, *Charadrius hiaticula* (Eur)
 Piping Plover, *Charadrius melodus* (NE)
 Mountain Plover, *Charadrius montanus* (GP)
 Eurasian Dotterel, *Charadrius morinellus* (Eur)
 Upland Sandpiper, *Bartramia longicauda* (NGP)
 Bristle-thighed Curlew, *Numenius tahitiensis* (AK)
 Curlew Sandpiper, *Calidris ferruginea* (Eur)
 Temminck's Stint, *Calidris temminckii* (Eur)
 Red-necked Stint, *Calidris ruficollis* (Eur)
 Little Stint, *Calidris minuta* (Eur)
 White-rumped Sandpiper, *Calidris fuscicollis* (A)
 Jack Snipe, *Lymnocyptes minimus* (Eur)
 Gray-tailed Tattler, *Tringa brevipes* (Eur)
 Spotted Redshank, *Tringa erythropus*. (Eur)
 Wood Sandpiper, *Tringa glareola* (Eur)
 Thick-billed Murre, *Uria lomvia* (NP)
 Long-billed Murrelet, *Brachyramphus perdix* (NWP)
 Kittlitz's Murrelet, *Brachyramphus brevirostris* (NP)
 Guadalupe Murrelet, *Synthliboramphus hypoleucus* (NP)
 Least Auklet, *Aethia pusilla* (NP)
 Swallow-tailed Gull, *Creagrus furcatus* (TP)
 Red-legged Kittiwake, *Rissa brevirostris* (NP)
 Ivory Gull, *Pagophila eburnea* (A)
 Black-headed Gull, *Chroicocephalus ridibundus* (Eur)
 Little Gull, *Hydrocoloeus minutus* (Eur)
 Ross's Gull, *Rhodostethia rosea* (A)
 Laughing Gull, *Leucophaeus atricilla* (TP)
 Black-tailed Gull, *Larus crassirostris* (NWP)
 Slaty-backed Gull, *Larus schistisagus* (Eur)
 Great Black-backed Gull, *Larus marinus* (NA)
 Least Tern, *Sternula antillarum* (TP)
 Red-billed Tropicbird, *Phaethon aethereus* (TP)
 Arctic Loon, *Gavia arctica* (Eur)

White-capped Albatross, *Thalassarche cauta* (SP)
 Wilson's Storm-Petrel, *Oceanites oceanicus* (Ant)
 Ashy Storm-Petrel, *Hydrobates homochroa* (EP)
 Providence Petrel, *Pterodroma solandri* (SP)
 Murphy's Petrel, *Pterodroma ultima* (SP)
 Hawaiian Petrel, *Pterodroma sandwichensis* (EP)
 Cook's Petrel, *Pterodroma cookii* (SP)
 Wedge-tailed Shearwater, *Ardenna pacifica* (TP)
 Great Shearwater, *Ardenna gravis* (SP)
 Magnificent Frigatebird, *Fregata magnificens* (TP)
 Blue-footed Booby, *Sula nebouxii* (TP)
 Snowy Egret, *Egretta thula* (S)
 Little Blue Heron, *Egretta caerulea* (S)
 Cattle Egret, *Bubulcus ibis* (S)
 Yellow-crowned Night-Heron, *Nyctanassa violacea* (S)
 White Ibis, *Eudocimus albus* (S)
 Glossy Ibis, *Plegadis falcinellus* (E)
 California Condor, *Gymnogyps californianus* (SW, 1897)
 Zone-tailed Hawk, *Buteo albonotatus* (SW)
 Yellow-bellied Sapsucker, *Sphyrapicus varius* (NE)
 Crested Caracara, *Caracara plancus* (SW)
 Eurasian Kestrel, *Falco tinnunculus* (Eur)
 Eurasian Hobby, *Falco subbuteo* (Eur)
 Dusky-capped Flycatcher, *Myiarchus tuberculifer* (SW)
 Variegated Flycatcher, *Empidonomus varius* (SA)
 Scissor-tailed Flycatcher, *Tyrannus forficatus* (SGP)
 Fork-tailed Flycatcher, *Tyrannus savana* (SA)
 Greater Pewee, *Contopus pertinax* (SW)
 Eastern Wood-Pewee, *Contopus virens* (E)
 Yellow-bellied Flycatcher, *Empidonax flaviventris* (NE)
 Alder Flycatcher, *Empidonax alnorum* (N)
 Eastern Phoebe, *Sayornis phoebe* (NE)
 Vermilion Flycatcher, *Pyrocephalus rubinus* (SW)
 White-eyed Vireo, *Vireo griseus* (SE)
 Bell's Vireo, *Vireo bellii* (E, SW)
 Yellow-throated Vireo, *Vireo flavifrons* (E)
 Blue-headed Vireo, *Vireo solitarius* (NE)
 Philadelphia Vireo, *Vireo philadelphicus* (E)
 Pinyon Jay, *Gymnorhinus cyanocephalus* (SW, 1967)
 Woodhouse's Scrub-Jay, *Aphelocoma woodhouseii* (SW)
 Eurasian Skylark, *Alauda arvensis* (Eur)
 Phainopepla, *Phainopepla nitens* (SW)
 Blue-gray Gnatcatcher, *Polioptila caerulea* (S)
 Brown Thrasher, *Toxostoma rufum* (E)
 Red-flanked Bluetail, *Tarsiger cyanurus* (Eur)
 Northern Wheatear, *Oenanthe oenanthe* (Eur)
 Dusky Thrush, *Turdus eunomus* (Eur)
 Redwing, *Turdus iliacus* (Eur)
 Siberian Accentor, *Prunella montanella* (Eur)
 Eastern Yellow Wagtail, *Motacilla tschutschensis* (Eur)
 Gray Wagtail, *Motacilla cinerea* (Eur)
 White Wagtail, *Motacilla alba* (Eur)
 Red-throated Pipit, *Anthus cervinus* (Eur)

Brambling, *Fringilla montifringilla* (Eur)
Hoary Redpoll, *Acanthis hornemanni* (N)
Lawrence's Goldfinch, *Spinus lawrencei* (SW)
Chestnut-collared Longspur, *Calcarius ornatus* (NGP)
Smith's Longspur, *Calcarius pictus* (A)
Thick-billed Longspur, *Rhynchophanes mccownii* (NC)
McKay's Bunting, *Plectrophenax hyperboreus* (AK)
Little Bunting, *Emberiza pusilla* (Eur)
Rustic Bunting, *Emberiza rustica* (Eur)
Field Sparrow, *Spizella pusilla* (E)
Lark Bunting, *Calamospiza melanocorys* (NGP)
LeConte's Sparrow, *Ammodramus leconteii* (NGP)
Nelson's Sparrow, *Ammodramus nelsoni* (NE)
Eastern Meadowlark, *Sturnella magna* (E)
Orchard Oriole, *Icterus spurius* (SE)
Hooded Oriole, *Icterus cucullatus* (SW)
Baltimore Oriole, *Icterus galbula* (NE)
Scott's Oriole, *Icterus parisorum* (SW)
Common Grackle, *Quiscalus quiscula* (NE)
Great-tailed Grackle, *Quiscalus mexicanus* (SW)
Ovenbird, *Seiurus aurocapilla* (NE)
Golden-winged Warbler, *Vermivora chrysoptera* (NE)
Blue-winged Warbler, *Vermivora cyanoptera* (NE)
Prothonotary Warbler, *Protonotaria citrea* (SE)
Lucy's Warbler, *Leiothlypis luciae* (SW)
Mourning Warbler, *Geothlypis philadelphia* (NE)
Kentucky Warbler, *Geothlypis formosa* (SE)
Hooded Warbler, *Setophaga citrina* (SE)
Cape May Warbler, *Setophaga tigrina* (NE)
Northern Parula, *Setophaga americana* (E)
Magnolia Warbler, *Setophaga magnolia* (NE)
Bay-breasted Warbler, *Setophaga castanea* (NE)
Blackburnian Warbler, *Setophaga fusca* (NE)
Blackpoll Warbler, *Setophaga striata* (NE)
Black-throated Blue Warbler, *Setophaga caerulescens* (NE)
Yellow-throated Warbler, *Setophaga dominica* (E)
Prairie Warbler, *Setophaga discolor* (E)
Black-throated Green Warbler, *Setophaga virens* (NE)
Canada Warbler, *Cardellina canadensis* (E)
Painted Redstart, *Myioborus pictus* (SW)
Summer Tanager, *Piranga rubra* (E)
Blue Grosbeak, *Passerina caerulea* (S)
Indigo Bunting, *Passerina cyanea* (NE)
Painted Bunting, *Passerina ciris* (SE)
Dickcissel, *Spiza americana* (GP)